Mon. Not. R. Astron. Soc. **000**, 000–000 (0000) Printed 27 November 2014 (MN LATEX style file v2.2)

Near-infrared studies of V5558 Sgr: an unusually slow nova with multiple outbursts

Ramkrishna Das[1] ⋆, Dipankar P. K. Banerjee[2] †, Arpita Nandi[1], N M Ashok[2] & Soumen Mondal[1]

[1] *Astrophysics & Cosmology Department, S N Bose National Centre for Basic Sciences, Salt Lake, Kolkata 700098, India.*
[2] *Astronomy & Astrophysics Division, Physical Research Laboratory, Navrangpura, Ahmedabad 380009, India.*

arXiv:1411.6524v2 [astro-ph.SR] 26 Nov 2014

27 November 2014

ABSTRACT

We present near-infrared (1-2.5 μm) JHK photo-spectroscopic results of the unusually slow nova V5558 Sgr (2007). V5558 Sgr showed a slow climb to maximum that lasted for about 60 days and then underwent at least five strong secondary outbursts. We have analyzed the optical light curve to derive large t_2 and t_3 values of 281 ± 3 and 473 ± 3 days respectively. An alternate approach is adopted to derive a distance estimate of 1.55 ± 0.25 kpc as conventional MMRD relation may not be applicable for a slow nova. In the pre-maxima stage the spectra showed narrow (FWHM ∼ 400 - 550 km s^{-1}) and strong emission lines of Paschen and Brackett series with prominent P-Cygni components. In the later phase the spectra show significant changes with the development of strong and broad (∼ 1000 km s^{-1}) emission lines of HI, HeI, OI, and NI and some uncommon Fe II emission lines. No evidence of dust formation is seen. V5558 Sgr has been shown to be a rare hybrid nova showing a transition from He/N to Fe II type from optical spectra. However the near-infrared data do not show such a transition and we discuss this anomalous behavior. A recombination analysis of the Brackett lines allows us to constrain the electron density and emission measure during the early optically thick phase and to estimate the mass of the ejecta to be (6.0 ± 1.5) × 10^{-4} M$_\odot$, assuming a filling factor of unity, from later observations.

Key words: infrared: spectra - line : identification - stars : novae, cataclysmic variables - stars : individual (V5558 Sgr) - techniques : spectroscopic, photometric.

1 INTRODUCTION

Nova V5558 Sgr was discovered by Y. Sakurai on 2007 April 14.777 UT (JD 2454205.277) at magnitude 10.3 (Nakano et al. 2007) during its pre-maximum stage. Henden & Munari (2007) have reported that no visible progenitor was detected in the POSS plates suggesting a value of 13.4 magnitudes as the lower limit of the outburst amplitude. They derived the astrometric position of the nova to be α = $18^h10^m18.258^s$ ± 0.046^s, δ = -18°46′51.95″ ± 0.047″ which is close to the coordinates reported by Sakurai (Nakano et al. 2007). After the discovery, the brightness of the nova increased very slowly. The maximum brightness was attained on July 10.0 UT at V = 6.53 (Munari et al., 2007b) which was followed by a flattening of the light curve and then a slow decline during which the nova underwent four more re-brightenings (Tanaka et al. 2011). Such a slow rise to maximum accompanied by multiple outbursts during the evolution have been rarely observed in other novae (for example, V723 Cas (Evans et al. 2003) and HR Del (Terzan 1970)).

Optical studies of V5558 Sgr have been performed by several observers (Iijima 2007a, 2007b; Naito 2007; Munari et al. 2007a; Poggiani 2008). The spectra recorded during the pre-maxima phase were dominated by the strong Balmer and He I lines along with weak features of emission lines of Fe II multiplets, Mg II, N II, Si II, Ca I and [O I]. All of these emission lines exhibited narrow and sharp profiles with typical full widths at half maxima (FWHM) of about 480-540 km s^{-1}, as measured from

⋆ E-mail: ramkrishna.das@bose.res.in
† E-mail: orion@prl.res.in

Balmer lines, which is a value range generally associated with Fe II type novae. Most of the strong emission lines were accompanied by absorption components of P-Cygni profiles, blueshifted by 400-500 km s^{-1}. Munari et al. (2007b) estimated a reddening of $E(B-V) = 0.36$ from the Na I D1 and D2 lines. No significant change in the spectra was noticed during the pre-maximum evolution except the disappearance of He I lines and the emergence of Fe II multiplets in emission on 2007 May 11 (Munari et al. 2007a).

Significant changes in the spectra were noticed during and after the initial maximum. The spectra, observed during and a short time after the initial maxima on 2007 July 10, were characterized by a smooth underlying continuum with numerous weak absorptions along with strong emission lines of Balmer series and FeII multiplets (Iijima 2007b, Munari et al. 2007b). Also, the He I lines which had disappeared on 2007 May 11, reappeared on 2007 July 12. Also a remarkable change was noticed in the line profile widths. The emission lines which were narrow in the pre-maxima stage, become strong and broad (FWHM $\sim$ 1150 km s^{-1}, Munari 2007b) which indicated the presence of fast hot winds. The nature of the spectra did not change much during the remaining part of the evolution till 2007 Nov 2 (Tanaka et al. 2011); the spectra were dominated by strong emission features of H, Fe II and He I lines along with weak emission features of OI, NII, CII. Many of these emission features were accompanied by sharp P-Cygni profile which disappeared gradually with time. Further spectroscopic follow up during the declining phase by Poggiani (2010) reported the appearance of forbidden lines, e.g., [OIII] 4959, 5007, and [Fe VI], [Fe VII], [Ca V] which suggests that V5558 Sgr had entered the nebular phase. Overall, the spectral behavior of V5558 Sgr before and after the maximum, is very similar to that of V723 Cas (Iijima 2007b; Poggiani, 2008, 2010). Poggiani (2010) estimated the characteristic time t_3 (time taken to decline by 3 magnitudes from visual maximum) = 170 $\pm$ 2 days making the nova a slow one according to the definition of speed class (Payne-Gaposchin 1957). Using the value of t_3 Poggiani (2010) revised the estimated value of the distance to be 1.3 - 1.6 kpc and absolute magnitude (M_V) at maximum in the range of 6.3 - 5.9 in agreement with the value of 6 $\pm$ 1 observed in case of slow novae V723 Cas and HR Del. Using these values Poggiani (2010) derived the white dwarf mass in the range of 0.58 - 0.63 $M_\odot$. We have certain reservations on Poggiani's (2010) method of deriving the distance which is discussed in section 3.2

The first infrared observations of V5558 Sgr were formally reported by Lynch et al. (2007). The infrared spectrum (0.8 - 2.5 μm) observed on 2007 September 12, showed strong and narrow (FWHM $\sim$ 620 km s^{-1}) emission lines of H I, He I, OI, Ca II, Fe II, and [N I]. All of the H I and few of the He I lines and Fe II lines showed P-Cygni absorption profiles which suggested significant optical depths in these lines. Further infrared observations performed by Rudy et al. (2007) on 2007 October 10 and 11 showed a spectrum (0.8 - 5.5 μm) dominated by low excitation broad (FWHM $\sim$ 1600 km s^{-1}) emission lines with strong N I and Fe II lines. The P-Cygni profiles had disappeared from the H I lines but were weakly present on the He I lines. They measured a reddening of about 0.8 from the O I lines. Both Infrared observations found no evidence of dust formation.

In this paper we present results of near-infrared JHK (1 - 2.5 μm) spectroscopic and photometric observations of V5558 Sgr. The observations are is described in Section 2 and the results in Section 3.

2 OBSERVATIONS

V5558 Sgr was observed in the near-Infrared JHK (1-2.5 μm) region using the 1.2m telescope at Mount Abu Infrared Observatory operated by Physical Research Laboratory, India. Observations were taken on 7 epochs in the pre-maxima rising phase and on 1 epoch in the post-maxima phase. The nova could not be followed in the months of mid-June to mid-October due to monsoon season. The near-infrared JHK spectra presented in this paper were obtained in each of J, H, K bands at similar dispersions of $\sim$ 9.75 Å/pixel using the Near Infrared Imager/Spectrometer with a 256×256 HgCdTe NICMOS3 array. Each time a set of at least two spectra of the object at two dithered position along the slit were taken. In the same manner, a set of two spectra of a nearby comparison star (SAO 161564, spectral type B9.2/A0V, $V = 5.126$) at similar airmass as the object were taken at the same grating position in order to remove the atmospheric absorption effects from the nova spectra. Each set of two images were subtracted from each other to eliminate the background counts which comprises of emission from sky and dark counts from the detector. From these subtracted images spectra were extracted using IRAF tasks. The spectra were wavelength calibrated using a combination of OH sky lines and telluric lines that register with the stellar spectra. Following the standard procedure, Hydrogen Paschen and Brackett absorption lines were removed from the spectra of the comparison star and the nova spectra were then divided by the spectra of the comparison star. The ratioed spectra were multiplied by a blackbody curve generated at the effective temperature of the comparison star to get the final spectra. However, caution should be taken while considering any spectral feature around 1.12 μm in the J band and between 2 to 2.05 μm in the K band, especially, in the early phase spectra when the emission lines are not so strong. This is because some differences in airmass between the nova and comparison star remained. Hence, the ratioing process, while removing telluric features, leaves some residuals in these wavelength regions where telluric absorption due to atmospheric oxygen and carbon-dioxide respectively are strong. The log of spectroscopic observations are provided in Table 1.

Near-Infrared photometric observations of the nova in the JHK bands were done in photometric sky conditions using the imaging mode of the NICMOS3 array. Several frames, both of the nova and a selected standard star (SAO 186544, spectral type B0.5Ib/IIC, $V = 6.02$) in each of the J, H, K filters, were recorded in 5 dithered positions offset typically by 20 arcsec. Near-infrared JHK magnitudes

Table 1. Observational log of the spectroscopic observations of nova V5558 Sgr. The date of outburst is taken to be 2007 Apr 14.777 UT.

Date 2007 (UT)	Days since Outburst	Integration time (sec) J	H	K
Apr. 26.951	12.174	90	90	40
Apr. 29.836	15.059	30	30	75
May. 01.861	17.084	60	60	40
May. 05.889	21.112	60	60	40
May. 07.832	23.055	30	30	90
Jun. 07.813	54.036	90	75	45
Jun. 10.844	57.067	75	75	40
Oct 20.594	188.817	60	60	90

were then extracted using APPHOT package in IRAF following the regular procedure for photometric reduction (e.g. Banerjee & Ashok, 2002) followed by us. The log of the photometric observations and the derived JHK magnitudes are given in Table 2. Further observations beyond November could not be carried out as the object had begun to approach solar conjunction.



Figure 1. A magnified view of the secondary outbursts in the optical and JHK light curves (black circles and gray crosses respectively) observed in V5558 Sgr. Optical data (V band and Visual) are taken from AAVSO and AFOEV databases. The near-Infrared JHK magnitudes are from the 1.2m Mount Abu Telescope (see Table 2) and from Walter et al. (2012). The errors of the optical and NIR data are typically smaller than 0.05 mag; smaller than their respective symbol sizes used.

3 RESULTS

3.1 Optical and near-infrared lightcurves

The partial V band light curve for the first 250 days, to specially highlight the secondary maxima, is shown in Figure 1 while the entire lightcurve over an extended period of 6.25 years from **maxima** is shown in Figure 2. In Figure 1 we also show the near-infrared JHK light curves. The optical light curve is based on the data collected from American Association of Variable Star Observers (AAVSO) and from Association Francaise des Observateurs dEtoiles Variables (AFOEV). The near-infrared JHK magnitudes are from Mt. Abu observations (see Table 2) and SMARTS/CTIO 1.5m telescope facility (Walter et al., 2012). The epochs of Mount Abu Infrared observations are marked by vertical dashes on the light curve. After the outburst the light curve rises very slowly. We find from AAVSO data that the nova took about sixty days to reach $V = 8.5$ from the discovery magnitude 10.3. After staying at $V = 8.5$, for a few days, the nova underwent its first episode of rebrightening to reach $m_v = 6.48$ on 2007 July 10.56 (JD = 2454292.06) which is in line with the findings of Munari et al. (2007b). Following this, the brightness of V5558 Sgr started to decrease. During the declining phase, V5558 Sgr showed at least four more secondary outbursts occurring on July 25 ($V = 7.7$, t = 101.6 days), August 12 ($V = 7.0$, t = 119 days), September 25 ($V = 7.1$, t = 163 days) and October 31 ($V = 7.4$, t = 199.6 days) respectively, where t = days after commencement of the eruption. After the fifth maxima V5558 Sgr declined slowly (see Figure 2). The near-infrared light curves are seen to mimic the behavior of the optical lightcurve and do not warrant any special comments. During the pre-maximum stage a gradual rising is seen in all the three near-infrared JHK bands similar to optical light curve, the J-H and H-K values were about 0.2 and 0.3 respectively. We do not have any near-infrared observation during the first maximum. However, the SMART/CTIO observations during the other secondary maxima show that the J-H and H-K have been about 1.1 and in the range of 0.6 - 1.0, respectively. From near-infrared observations we do not notice any excess in any of the near-infrared JHK light curves which indicates that no dust has been formed in the nova ejecta.

The light curve shows all the characteristics of the J class of novae based on the classification system of Strope et al. (2010). J class of nova show 'jittering' or 'flare-ups' of substantial amplitude which could be thought of as outbursts above a smoothed light curve that follows the universal decline law predicted by Hachisu & Kato (2006). Strope et al. (2010) listed 14 novae that belong to J type. While we compare the light curve of V5558 Sgr with these novae, we find close similarities with V723 Cas and HR Del out of these 14 novae. Both of V723 Cas and HR Del showed a prolonged pre-maximum halts, V723 Cas took 100 days and HR Del took 180 days respectively to reach the maximum. Both of these novae showed multiple outbursts, V723 Cas showed 6 outbursts in between 100 and 425 days and HR Del showed 5 outbursts in between 180 and 380 days.

The cause behind the occurrence of multi-outbursts, as observed in this nova, is not well understood. Previous studies of slow novae have shown that multi-outbursts might take place on white dwarfs of mass ~ 0.6 $M_\odot$ which is close to the critical value required beginning of the thermonuclear runaway event (for example, Friedjung et al. 1992; Kato et al. 2002). However, recent calculations with OPAL opacities,

Table 2. Observational log of the photometry. The date of outburst commencement is taken to be 2007 Apr 14.777 UT.

Date 2007 (UT)	Days since Outburst	Magnitudes J	H	K
Apr. 29.891	15.114	7.02 ± 0.05	6.57 ± 0.09	6.09 ± 0.05
May 01.955	17.178	6.94 ± 0.06	6.71 ± 0.10	6.03 ± 0.10
May 05.963	21.186	6.77 ± 0.11	6.35 ± 0.07	5.93 ± 0.13
May 07.944	23.167	6.70 ± 0.09	6.33 ± 0.05	5.89 ± 0.07
Jun. 07.860	54.083	6.29 ± 0.12	5.80 ± 0.08	5.59 ± 0.12
Jun. 10.798	57.021	6.25 ± 0.10	5.84 ± 0.12	5.48 ± 0.09

show that thermonuclear runaway may occur even in much less massive white dwarfs (WDs), e.g., 0.2 - 0.5 M⊙ WDs (Shen et al. 2009). In a recent study, Kato & Hachisu (2009, 2011) pointed out that two different types of evolution of slow novae of a certain range of WD masses, ∼ 0.5 - 0.7 M⊙ may take place. One is the evolution with optically thick winds and the other is without optically thick winds. The wind-type novae show a sharp peak in the optical light curve and the light curve decays quickly. On the other hand novae without an optically thick wind evolve very slowly and stay at an extended low-temperature stage for a long time, which makes a long-lasting flat optical peak. They suggested that transition from the state of non-optically thick winds to a state of optically thick winds may take place. Such a transition may be accompanied by violent activities like oscillatory behaviors in the light curves of few slow novae which are considered as some relaxation processes associated with the transition. This kind of transition in low-mass WDs (⩽ 0.6 M⊙) in close binary system, may be triggered by the companion star through the effects of (1) spin-up by the companion motion, (2) gravity of the companion star, and (3) drag luminosity due to frictional energy deposition. Incorporating these effects, the authors **(Kato & Hachisu 2011)** generated two trial models adopting a parameter set of the mass of white dwarf and companion, chemical composition of the envelope and the orbit. The trial light curves successfully reproduced the flat phase of static evolution followed by a smooth decline due to wind mass loss. From this study the WD mass was estimated to be about 0.6 M⊙ for all three novae, V723 Cas, HR Del, and V5558 Sgr.



Figure 2. The V band optical light curve of V5558 Sgr, plotted for ∼ 6.25 years after outburst, is shown, in the top panel. The data are restricted to either photoelectric or CCD measurements from the AAVSO and AFOEV databases. The grey curve shows that the light curve can be well matched by a uniform curve, obtained from a polynomial fit, in case the data points for the secondary maxima are excluded (see text for additional details). The bottom panel shows the plot of the extinction A_k in the K band versus distance towards the nova (filled dark circles; from Marshall et al. 2006; $A_v = 11A_k$). The intersection point of this extinction curve with the MMRD generated curve can be used to determine the distance and extinction to the nova simultaneously. See text for more details.

3.2 Distance to the nova

If the data points of the 5 major rebrightenings or secondary maxima are excluded, then it is found that the entire light curve can be fit by one uniform curve that initially rises and then declines. This uniform curve, on which the secondary maxima sit, is shown by the gray line which is obtained by a least-squares minimized polynomial fit. From this smooth curve, we estimate that maximum to be reached on 5 July 2007 at V = 8.3; t_2 and t_3, the times to decline by 2 and 3 magnitudes from visual maximum, to be 281 ± 3 and 473 ± 3 days respectively. V5558 is therefore an unusually slow nova; possibly one of the slowest ever known (from the selection of well-observed novae by Warner(1995), only DO Aql has larger t_2 and t_3 values of 430 and 900d respectively). The validity of using maximum magnitude - rate of

decline (MMRD) relations for such extremely slow novae is debatable because most MMRD relations are obtained as outcomes of fits of M_v to t_2 or t_3 values in which the latter have much smaller values. For example, considering two well known relations, in the sample used by della Valle & Livio (1995) the largest value of t_2 is 117d and in the Downes and Duerbeck (2000) MMRD sample there is only one nova with t_2 greater than 100 days (viz. HR Del with t_2 = 172d). In general not many very slow novae are known; Capaccioli et al. (1990) cite the case of one nova in the LMC with t_2 = 117d while Darnley et al (2006) give instances of four novae with t_2 exceeding 100d viz. 817.2, 198, 100.3 and 331d respectively.

We thus estimate M_v and the distance to the nova using a different approach. The variation of the extinction with distance towards the nova is shown, for a ten arc minute field around the object, in the bottom panel of Figure 2 based on modeling of the galactic extinction by Marshall et al. (2006). Two values of the excess $E(B-V)$ have been estimated for V5558 Sgr using independent approaches. From the interstellar NaI D1 and D2 lines, Munari et al (2007a) obtain $E(B-V) = 0.36$ while Rudy et al (2007) obtain $E(B-V) \sim 0.8$ based on the relative strengths of the O I 8446 and 12187Å lines. Both methods are known to yield reliable estimates of the reddening. Using both these values of $E(B-V)$, and the relations $A_v = 3.1E(B-V)$ and $A_v = 11A_k$, the distance to the nova from Figure 2 (bottom panel) is then constrained to lie in the range 1.3 and 1.8 kpc (i.e $d = 1.55 \pm 0.25$ kpc).

3.3 Line identification and general characteristics of the *JHK* spectra,

The near-infrared *JHK* spectra of V5558 Sgr are presented in Figures 3, 4 and 5, respectively. The lines are marked on the figures and a list of the identified lines are given in Table 3. The epochs of our observations are indicated by bars on the abscissa on top in Figure 1. The presented near-infrared data covers the premaxima evolution till Jun 10, our last data is on Oct 20, 11 days before the fifth maximum which was reached on 31 October. None of our spectra unfortunately coincide with the rebrightenings to monitor spectral changes that took place at these secondary maxima.

In the initial stage, during 2007 April 26 - June 10, the *JHK* spectra show strong emission lines of HI; these include Paγ and Paβ in the J band, Br10 to 19 in the H band and Brγ in K band. Except the Paschen and Brackett series lines, the only other discernible feature is the NI 1.2461, 1.2469 μm line which becomes much stronger in the later phase on 2007 Oct 20. In the *J* band spectra OI lines at 1.1287 and 1.3164 μm which are generally seen in classically novae spectra, are weak or not present. But it must be noted that the OI 1.1287 micron line lies in a region of poor atmospheric transmission and so it may be weakly present but not clearly discernible. An important feature to be noted is all the emission lines are very narrow. In order to quantify the changes in the linewidths, the evolution of Pa β and Brackett γ lines which are prominent but not blended with any other line, was investigated. The full width at half maximum (FWHM) of the Pa β and Br γ profiles are estimated to be 533 km s^{-1} and 387 km s^{-1} respectively. Correcting these for instrumental broadening, our *J* band has poorer resolution than the *Kband*, gives values of $\sim$ 240 - 260 km s^{-1} for both lines indicating a very slow velocity for the ejecta. During the pre-maxima phase, there were no significant changes seen in the width of the emission lines implying the absence of any significant change in the expansion velocity of the ejected envelope. During the course of pre-maximum evolution we also note that all the emission lines show sharp P-Cygni absorption features on the blue wings. The values of these absorption components for the Br 12 and Brackett γ lines (Figure 6) are measured approximately at 530 km s^{-1} from the emission peak.

Significant changes in the spectra are observed on 2007 Oct 20, a few days before the fifth brightening. The nova had entered the nebular stage at this time (Poggiani 2010; Tanaka et al. 2011). The *JHK* spectra show strong emission lines of Paschen and Brackett series, HeI, NI, OI and Fe II. Among the He I lines, those at 1.0830, 1.7002 and 2.0581 μm are prominent. The Lyα fluoresced OI line 1.1287 μm and continuum fluoresced OI line 1.3164 μm, which were absent in the pre-maximum phase, are present as normally seen in the spectra of classical novae. The large observed ratio ($\sim$ 2.7) of the OI 1.1287/1.3164 μm lines indicates that Lyα fluorescence is the dominant process contributing to the strength of the 1.1287 μm line. In *J* band, a magnified view of the region between 1.19 and 1.26 μm shows a complex blend of several weak but well resolved emission lines. NI is clearly detected as a broad feature centered around 1.246 μm (the NI 1.2461 and 1.2470 μm lines) on 2007 Oct 20. Other features at 1.998, 1.2086, 1.2211, 1.2305, 1.2581 μm, in this region are also detected which we attribute to NI lines. Few of these features are possibly blended with adjacent weak FeII lines (see Table 3 for details). In addition, two strong features at 1.3448 in *J* band and 1.7878 μm in *H* band are detected which are assigned to NI.

In addition to the HI, HeI, OI and NI lines, we see prominent emission features of FeII in each of the *JHK* bands also. These are at 1.1126 μm which is one of the so-called '1 micron Fe II lines' (Rudy et al. (2000)) in *J* band, 1.6872 and 1.7414 μm in *H* band. All of these Fe II lines are rarely observed in the spectra of novae, for example, in V2615 Oph (Das et al. 2009), RS Oph (Banerjee et al. 2009, V574 Pup (Naik et al. 2010) and V5588 Sgr (Munari et al 2014; in preparation). Apart from these, there are two more novae, namely V2540 Oph (Rudy et al. 2002) and C I Aql (Lynch et al. 2004), where these lines appear to be detected. The excitation mechanism for these lines is believed to be Lyman α and Lyman continuum fluorescence coupled with collisional excitation (Banerjee et al. 2009 and references therein). Further, strong emissions of these Fe II lines suggest high density conditions (10^{10} - 10^{12} cm^{-3}) inside the ejecta which is supported by recombination analysis (section 3.6) also. As the detections of these Fe II lines in individual objects increase, it becomes amply evident that these *H* band lines could be present in the spectra of other novae too, but have evaded detection because of blending - especially when their widths are large - with the Br 11 (1.6806 μm) line. In FeII class of novae additional blending will take place



Figure 3. The J band spectra of V5558 Sgr on different days with the flux normalized to unity at 1.25 μm. The prominent lines are identified and marked (see Table 3 for details; u.i = unidentified)

with the strong CI line at 1.6890 μm and also CI 1.7449 μm . An instance where these FeII lines are very clearly resolved is in RS Ophiuchi (Banerjee et al. 2009), especially during the later stages of its outburst when all line widths in RS Oph narrowed due to deceleration of the ejecta.

We also detect weak emission features at 2.1140 and 2.1470 μm. The 2.1140 μm feature which was observed in V723 Cas also, is quite certainly a blend of HeI 2.1120, 2.1132 lines. The line at 2.1441 μm is probably due to Fe II. Two possible identifications may be considered for the emission feature at $\sim$ 2.090 μm in the K band. First, it could be FeII 2.0888 μm, a line also seen in the nova V2615 Oph (Das et al. 2009)for which an excitation mechanism by Lyman α fluorescence was proposed. Alternatively, this feature could be the [Mn XIV] 2.0894 μm coronal line which has been seen in a few instances in novae spectra during the coronal phase viz., in nova V1974 Cyg (Wagner & Depoy, 1996) and in RS Oph (Banerjee et al. 2009). However a lack of detection of optical coronal lines at this time (Poggiani 2010) or of any other coronal lines in our spectra is not supportive of a [Mn XIV] assignment. The present 2.0888 μm FeII line should not be confused with an unidentified line at 2.0996 μm that has often been detected in novae and which still remains unidentified (Rudy et al. 2002)

The emission lines observed on 2007 October 20 were significantly broader in comparison to the pre-maximum stage. We measured FWHM as 1287 km s^{-1} from Pa β, and 886 km s^{-1} from Br γ lines (instrumental deconvolved widths of 1201 and 880 km s^{-1} respectively . While this result matches with the findings of Munari et al. (2007b)it is puzzling why two lines from the same species should have significantly differing velocities It is possible that they arise from different regions within the ejecta; this is known to happen in the ionized plasma discs of Be stars. No strong absorption feature associated with the emission lines is noticed. It is also noted that no lines from low ionization species such as Na I or MgI are seen in the JHK spectra. These low ionization lines, which are indicative of low temperature conditions, have been suggested as potential diagnostic features to predict dust formation in the nova ejecta (Das et al. 2008). The absence of these lines in V5558 Sgr is consistent with the lack of dust formation in this nova (Lynch et al. (2007), Rudy et al. (2007)).



Figure 4. The *H* band spectra of V5558 Sgr on different days with the flux normalized to unity at 1.65 μm. The prominent lines are identified and marked (see Table 3 for details.)

3.4 Spectral classification

Williams (1992) introduced two principal classes for novae, the "Fe II" and "He/N" classes, based on the strength of of non-hydrogen emission lines in the optical spectra in the early stage. In addition, a small fraction of novae display characteristics of both classes and these are referred to as "hybrid" or "Fe IIb" novae. These novae have strong Fe II lines soon after the outburst which are broader than those seen in typical "Fe II" novae and subsequently display "He/N" spectrum. An example of "hybrid" nova is V574 Pup (Naik et al. 2010; also see Williams 1992). The optical classification scheme has been extended into near-Infrared spectral region *JHK* (1.08 to 2.35 μm) by Banerjee & Ashok (2012). In the near-infrared, the principal differentiating feature between the two classes is the strong presence of carbon lines in "Fe II" novae and their absence in He/N novae. The strongest of these C lines appear in the J band at 1.166 and 1.175 μm and in the H band at 1.6890 μm and several lines between 1.72 to 1.79 μm. On the other hand, He/N novae show prominent He emission right from the beginning of the outburst. The principal He lines are generally seen at 1.0830, 1.7002, 2.0581 and the 2.1120,2.1132 μm. Few near-infrared lines of nitrogen are detected in both Fe II or He/N spectra (for e.g the 1.2461, 1.2469 μm lines) but they are invariably stronger in He/N novae as observed in optical spectra also (Williams 1992).

V5558 Sgr comes closest to being classified as a hybrid nova but with an anomalous component. The hybrid tag is justified because it showed a transition from one class to another but the anomaly arises because the transition was in the reverse direction viz from He/N to FeII instead of the other way around. Only one other hybrid nova T Pyx has shown such a similar reverse transition, both in the optical and near-infrared (see Joshi et al. 2014 for the near-infrared behavior and references therein for the optical results). Let us review the optical and near-infrared behavior of V5558 Sgr and compare them using the present results and those from Tanaka et al (2011) because they share common dates (or very nearby) of observation between the two.

In the optical, on April 16 it appeared as a He/N type nova but by 2007 April 25 Fe II lines and a Fe II class spectrum appeared showing that it had evolved from the He/N type to the Fe II type. T Pyx made a similar transition within a week or less (Joshi et al. 2014). In the near-infrared, there is no spectrum on 16 April to compare with the optical. But for the Tanaka et al (2011) spectrum of 25 April in the optical, the near-infrared spectra of 26 April is available and



Figure 5. The K band spectra of V5558 Sgr on different days with the flux normalized to unity at 2.20 μm. The prominent lines are identified and marked (see Table 3 for details.).

this is certainly not of the Fe II type. No prominent CI lines are present as was clearly seen in T Pyx. The near-infrared spectra at this stage more closely resemble the He/N class (there is a caveat attached to this sentence which we qualify later). All subsequent optical spectra between 25 April to 10 June are of typical of the Fe II class but surprisingly all the near-infrared spectra during this same time are more close to the He/N type. This is a major difference in the optical and **infrared** behavior and a significant finding from this work. This adds complexity to any unification scheme (Williams 2012; Shore 2013) that tries to explain Fe II, He/N and hybrid novae simultaneously.

In the paradigm proposed by Williams (2012), the the He/N spectra have their origin in the material lost from the white dwarf surface during the thermonuclear runaway. That is why the He lines, responsible jointly with N lines for the creation of the He/N tag, are only seen immediately after the outburst before the He-rich nova ejecta material intermingles with a large circumbinary envelope of gas whose origin is the secondary star. The dominance of the He lines is thus decreased. The Fe II spectra point to their formation in this large circumbinary envelope of gas which is proposed to be created by the impact of the nova ejecta with the secondary star. Depending on the proximity of the secondary star, the WD ejecta mass, and the radiation field impinging on the secondary star, the secondary star could be stimulated to eject mass. Hybrid objects are explained by changing parameters in the two emitting regions (one comprising of the WD ejecta material and the other of the circumbinary envelope) during the post outburst decline. How the contrasting IR behavior (vis-a-vis the optical behavior) in V5558 Sgr could be reconciled within this scheme will be interesting to see. We do not have a easy or straightforward answer to explain the observed behavior but are working on a general model that tries to reproduce both the Fe II and He/N type spectra. In such a model, we feel there is a need for including an additional parameter which considers the changing excitation and ionization conditions in the ejecta resulting from changes in the central WD's temperature.

If we do not take help of the optical spectra and base a classification solely on just the near-infrared spectra during and around maximum (i.e. spectra between April to June), the caveats associated with a He/N classification throughout this period are as follows. First the He lines, the important ones being expected at 1.0830, 1.7002 and 2.0581, are very weak or missing. However NI is present in the form of NI 1.2461, 1.2469 μm right from the earliest stage at a strength roughly consistent with that expected in the He/N class. But

Table 3. List of observed lines along with fluxes in the JHK spectra.

Wavelength (μm)	Species	Other contributing lines	Line fluxes (in 10^{-15} W cm^{-2} μ^{-1}) Apr 29	May 05	Jun 10	Oct 20
1.0830	He I		...	...	...	973.0 ± 27.0
1.0938	Pa γ		32.7 ± 8.0	50.7 ± 3.0	72.3 ± 5.0	1240.0 ± 30.0
1.1121	u.i. [a]/ Fe II?		...	...	...	80.5 ± 5.0
1.1287	O I		...	...	...	221.0 ± 20.0
1.1998	N I		...	...	...	21.4 ± 1.4
1.2087	N I		...	...	...	18.5 ± 0.5
1.2211	N I		...	...	...	10.0 ± 1.0
1.2305	N I		...	...	...	16.0 ± 0.5
1.2393	u.i. [a]/Fe II?		...	...	...	7.0 ± 0.5
1.2461,70	N I		3.2 ± 0.7	3.7 ± 0.5	0.8 ± 0.2	45.0 ± 4.0
1.2581	N I		...	...	...	17.0 ± 0.5
1.2818	Pa β		67.8 ± 5.0	57.8 ± 3.5	66.7 ± 4.0	985.0 ± 17.0
1.3164	O I		...	...	...	80.5 ± 2.0
1.3448	N I		...	...	...	32.0 ± 0.5
1.5260	Br 19		2.5 ± 0.3	2.0 ± 0.4	1.4 ± 0.3	13.4 ± 2.0
1.5342	Br 18		3.1 ± 0.5	2.3 ± 0.3	1.9 ± 0.4	20.9 ± 2.0
1.5439	Br 17		3.1 ± 0.4	2.4 ± 0.5	1.7 ± 0.4	25.2 ± 3.0
1.5556	Br 16		4.1 ± 0.5	3.5 ± 0.3	2.8 ± 0.3	78.7 ± 4.0
1.5701	Br 15	Fe II 1.5748?	2.3 ± 0.3	2.3 ± 0.4	1.9 ± 0.5	84.1± 4.0
1.5881	Br 14		4.4 ± 0.4	3.2 ± 0.3	3.0 ± 0.4	70.0 ± 3.0
1.6109	Br 13		4.2 ± 0.3	3.6 ± 0.4	3.1 ± 0.3	91.0 ± 3.0
1.6407	Br 12		4.2 ± 0.3	3.7 ± 0.4	2.8 ± 0.4	120.0 ± 5.0
1.6806	Br 11	Fe II 1.6872	4.9 ± 0.3	5.4 ± 0.6	4.2 ± 0.5	166.6 ± 5.2
1.6872	Fe II		0.4 ± 0.1	0.5 ± 0.1	0.8 ± 0.3	142.0 ± 5.5
1.7002	He I		...	...	...	28.8 ± 2.0
1.7362	Br 10	Fe II 1.7414	6.4 ± 0.3	6.1 ± 0.5	5.5 ± 0.4	205.1 ± 4.8
1.7878	N I		...	...	...	16.7 ±2.5
2.0581	He I		...	...	...	420.0 ± 6.0
2.0888	Fe II	[MnXIV] 2.089?	...	...	...	26.5 ± 0.2
2.1120,32	He II		...	...	...	14.0 ± 0.2
2.1470	Fe II		...	...	...	7.6 ± 1.0
2.1655	Br γ		6.8 ± 0.3	7.6 ± 0.4	7.2 ± 0.4	410.0 ± 6.0

[a] u.i. = unidentified

there is another troubling anomaly. The observed width of the lines are very narrow whereas He/N novae invariably have line widths of several thousands of km/s. Thus, on the whole, the spectra of V5558 Sgr emphasizes both the complexity and the diversity of novae spectra.

3.5 Recombination analysis

A recombination case B analysis of the HI line strengths was performed for the spectra of all days. Representative results for four days viz. 2007 April 29, May 05, Jun 10 and Oct 20 are presented in Figure 7 which represents the observed strength of the Brackett lines with respect to Br14 normalized to unity. Debelending of the H lines with contaminating lines (for e.g. Br 10 and Br 11 with FeII lines) was done either manually (see Figure 8 as an example) or using IRAF tasks. The observed H line strengths, presented in Table 3, were then compared with that expected by using case B emissivities from Storey & Hummer (1995). From recombination theory it is qualitatively expected, under optically thin conditions, that when strengths of lines of the same series are compared, a lower line of the series should be stronger than a higher line. For example, it is expected



Figure 6. The time evolution of the Br12 and Br γ 2.1655 μm lines showing the P-Cygni profile in the early stages with its absorption feature at $\sim$ 530 km s^{-1} blueward of the emission peak.

that Brγ (corresponding to a transition between levels 7–4) is expected to be significantly stronger than any higher line of the series like Br10 or Br11 (transitions between 10 – 4 and 11 – 4 respectively). But the reverse is actually being observed here on all epochs except for 2007 20 October. This clearly indicates that at these epochs the Br γ line is optically thick and possibly so are the other Br lines also. The comparison has been made here with a representative Case B model at T = 10000K and density = 10^{10} cm^{-3} but our conclusions will remain equally valid for case B values over an extended parameter space of electron density n$_e$ varying between 10^5 and 10^{13} cm^{-3} and temperature varying between 5000 and 20000 K. This may be seen for e.g. in a similar detailed graphical illustration done for the novae RS Oph **(Figure 6, Banerjee et al. 2009)** and T Pyx **(Figure 9, Joshi et al. 2014)**.

Because the Br lines appear optically thick, their actual strengths are unknown and hence it is not possible to estimate the mass of the nova by comparing observed and predicted case B line luminosities. However, the emission measure $n_e{}^2L$ can be constrained where L is the linear size of the emitting region. The values of the opacity $\Omega_{n,n'}$ for different HI lines for transitions between levels (n, n') at different temperatures and densities are known (tabulated in Hummer & Storey 1987; Storey & Hummer 1995). The optical depth at line-center $\tau_{n,n'}$ can be calculated using $\tau = n_e n_i \Omega L$, where L is the path length in cm. From Figure 8, among the Brackett lines, it can be clearly seen the Br γ line is definitely optically thick. Thus we use the Brγ line and assume with considerable certainty that its optical depth is greater than unity i.e. τ(Brγ)$>$ 1.

From Storey & Hummer (1995) we note that the value of the opacity factor $\Omega_{n,n'}$ does not vary too much with density or temperature for the near-infared HI lines including Brγ (the quantitative change is mentioned a sentence later). Since in the early outburst stages, the electron density in the nova ejecta is expected to be high (values as high as $n_e = 10^{14}$ cm^{-3} are even invoked) we make a conservative estimate that n_e is in the range 10^9 to 10^{13} cm^{-3}. Using the range of values of $\Omega_{n,n'}$ between 1.3 $\times$ 10^{-34} to 7.46 $\times$ 10^{-34} corresponding to n_e values varying between 10^9 to 10^{13} cm^{-3} for a representative temperature $T_e = 10^4$ K, we get the emission measure $n_e{}^2L$ to be in the range 1.3 $\times$ 10^{33} to 7.7 $\times$ 10^{33} cm^{-5}.

Approximate constraints on the electron density can be obtained if we can take L to be the kinematic distance $v\times t$ traveled by the ejecta where v is the velocity of ejecta and t is time after outburst. Using a representative value of v = 250 kms^{-1} as discussed earlier and t ranging from 12 to 188 days, and applying the constraint that τ(Brγ)= $n_e n_i \Omega L >$ 1; we obtain lower limits for n_e in the range 1.7 $\times$ 10^9 cm^{-3} to 1.6 $\times$ 10^{10} cm^{-3} (assuming $n_e = n_i$). These lower limits are likely to be smaller than the actual n_e values because τ(Brγ) can be considerably $>$ 1 that has been used above. The density in the ejecta is seen to be fairly high over the duration of our observations. It is worth noting Lynch et al. (2000) show that high densities of 10^{10} cm^{-3} or more tend to thermalize the level populations through collisions and thereby bring about deviations from Case B predictions as is observed here.

By 20 October, the ejecta appears to have diluted sufficiently to be optically thin and the line strengths follow a caseB distribution well. We estimate of the mass of the emitting gas at this epoch using the same approach as in some earlier novae by using the following relation (Banerjee et al. 2010, Raj et al. 2013):



Figure 7. Recombination analysis for the hydrogen Brackett lines in V5558 Sgr on selected dates. The abscissa is the upper level number of the Brackett series line transition (Br7 = Brγ). The line intensities (filled circles) are relative to that of Br 14. The Case B model predictions for the line strengths are also shown by dashed lines.

$$M = (4\pi d^2 (m_H)^2 (fV/\epsilon))^{0.5} \quad (1)$$

where d is the distance, m_H the proton mass, f the observed flux in a particular line, ϵ the corresponding case B emissivity; V is the volume of the emitting gas which equals $[4/3\pi(vt)^3\phi]$, where ϕ, v and t are the filling factor, velocity and time after outburst, respectively. We assume T = 10000K and n$_e$ to be $\sim$ 10^{10} cm^{-3} prompted both by the goodness of the fit in the bottom panel of Figure 7 and for the following additional reason. As discussed above, n$_e$ was close to $\sim$ 10^{10} cm^{-3} during our June observation (57d after outburst) but most likely had an even higher value. If the ejecta subsequently diluted geometrically (n$_e$ $\propto$ r^{-2}) thereafter upto 20 October (188d after outburst), the density would drop by a factor of 10 (i.e. square of 57/188). Based on this choice of n$_e$, we estimate the mass of the gas in the ejecta in the range (6.0 $\pm$ 1.5) $\times$ 10^{-4} M$_\odot$. This is a standard value for novae. However, this mass estimate is restricted in accuracy by our lack of knowledge of the filling factor which is assumed to be 1 here .



Figure 8. Deblending of the 2007 Oct 20 profiles of the Br 11 (1.6806 μm) and B10 (1.7362 μm) lines with neighboring Fe II 1.6872 and 1.7414 μm lines respectively using two-gaussian fits. The individual gaussians are shown by the dotted lines, their co-added sum by the dashed lines and the observed data by the gray lines.

3.6 Evolution of the continuum

We analyze the evolution of the continuum flux distribution to check for any sign of dust formation and also analyze the general trend in the temporal evolution of the spectral energy distribution (SED). The continuum has been modeled with power-law fits ($F_\lambda = \lambda^{-\alpha}$), the results of which are shown in Figure 9, wherein representative spectra on 2007 April 29, May 5, June 10 and October 20, sampling the duration of our observations have been plotted. The spectra have been flux calibrated using contemporaneous JHK values from Table 2 or data from Walter et al. (2012) after correcting these for reddening using E(B - V) = 0.58 as the average of the values provided by Munari et al. (2007a) and Rudy et al. (2007). In the early stage i.e. during April 26 - May 5 of the evolution of V5558 Sgr, the continuum spectrum is well matched with fits having spectral indices of $\alpha \sim 1.85$ - 1.9. The subsequent spectra, observed during Jun 7 - 10, tend to become slightly steeper with the slope of $\alpha = 2.1$. Around the fifth maximum the SED becomes flatter with $\alpha = 1.4 \pm 0.5$. Proceeding further, we have also specifically tried to fit the SED for 2009 February 25 which is about 317 days after the discovery. The result is shown in the last panel in Figure 9 where we have used JHK data from Walter et al. (2012) and BVI magnitudes from AAVSO database. We find that the continuum is well fitted by a power law, with the spectral index 1.95 ± 0.5.

Such study of evolution of continuum has been performed in case of some other novae also. For example, in V574 Pup (Naik et al. 2010) approximate value of the spectral index α was observed to decline from a value of 2.75 in the initial stage. In KT eridani (Raj et al. 2013) the value of the spectral index was found to decrease from 3.34 to 2.67. However, in the case of V4633 Sgr (Lynch et al. 2001) the change in the slope of the continuum was found to be in the opposite direction. The slope changed from 2 to 2.7 in observations taken 525 and 850 days after outburst. In comparison, the behavior of V5558 Sgr is peculiar where α did not increase or decrease monotonically as observed in other novae. This appears to be a consequence of the very slow evolution of the nova. It may be noted that a blackbody, in the Rayleigh-Jeans regime, is expected to give a index of 4, quite different from that seen here.



Figure 9. The SED for nova V5558 Sgr is shown using $BVIJHK$ fluxes for five representative dates 2007 April 29, 2007 May 5, 2007 June 10, 2007 October 20 and 2009 February 25 are shown. BVI fluxes are from AAVSO, JHK fluxes on first four days are from Mount Abu observations. JHK magnitudes on 2009 February 25 are collected from SMART observations. The composite JHK spectra for the first four representative dates are shown also. Model fits to the data with a power law, $F_\lambda \propto \lambda^{-\alpha}$, are shown by the continuous lines with the broad-band fluxes represented as filled circles.

No evidence for dust formation is seen, which in case it is formed, will show up as a significant excess at the infrared wavelengths. This is not the case here. Observations by the Wide field Infrared Survey Explorer (WISE) too, at an even later date on 2010 23 March, do not show any source with infrared excess at the position of the nova indicating the non-formation of dust.

4 SUMMARY

We have presented multi-epoch near-infrared spectroscopy and photometry observations of the slow nova V5558 Sgr which was discovered in April 2007 and which showed a sequence of multiple secondary outbursts. The optical light curve has been analyzed allowing a distance estimate of 1.55 $\pm$ 0.25 kpc to be made. V5558 Sgr is a fairly unique nova by virtue of showing a rare reverse-hybrid transition from He/N to Fe II type, based on its optical spectra. However the near-infrared data do not show such a transition and a discussion is made of this aspect. A Case B recombination analysis of the hydrogen lines has been done allowing the mass of the gaseous component in the ejecta to be estimated to be (6.0 $\pm$ 1.5) $\times$ 10^{-4} $M_{\odot}$ assuming a filling factor of unity.

5 ACKNOWLEDGMENTS

The research work at S N Bose National Centre for Basic Sciences is funded by the Department of Space and Technology, Government of India. The research work at Physical Research Laboratory is funded by the Department of Space, Government of India. We are thankful to AAVSO, USA and AFOEV, France for the use of their optical photometric data.

REFERENCES

Banerjee D. P. K., Ashok N. M., 2012, BASI, 40, 243
Banerjee D. P. K., Ashok N. M., 2002, A&A, 395, 161
Banerjee D. P. K., Das R. K., Ashok N. M., 2009, MNRAS, 399, 357
Banerjee D. P. K., Das R. K., Ashok N. M. et al., 2010, MNRAS, 408, L71
Capaccioli M., Della Valle M., D'Onofrio M. & Rosino L., 1990, 360, 63
Darnley M. J., Bode M. F., Kerins E. et al., 2006, MNRAS, 369, 257
Das R.K., Banerjee D. P. K., Ashok N. M., Chesneau O., 2008, MNRAS, 391, 1874
Das R.K., Banerjee D.P.K., Ashok N.M., 2009, MNRAS, 398, 375
della Valle M., Livio M., 1995, ApJ, 452, 704
Downess R. A., Duerbeck H. W., 2000, AJ, 120, 2007
Evans A., Gehrz R. D., Geballe T. R. et al., 2003, AJ, 126, 1981
Friedjung M., 1992, A&A, 262, 487
Henden A., Munari U., 2007, IBVS, 5803
Hummer D. G., Storey P. J., 1987, MNRAS, 224, 801.
Iijima T., 2007a, CBET, 934
Iijima T., Correia A. P., Hornoch K., Carvajal J., 2007b, CBET, 1006
Joshi V., Banerjee D. P. K., Ashok N. M., 2014, MNRAS, 443, 559
Kato T., Uemura M., Haseda K., Yamaoka H., Takamizawa K., Fujii M., Kiyota S., 2002, PASJ, 54, 1009
Kato, M., Hachisu, I. 2009, ApJ, 699, 1293
Kato, M., Hachisu, I. 2011, ApJ, 743, 157
Lynch D.K., Wilson J. C., Rudy R. J., Venturini C. C., Mazuk S., Miler N. A., Puetter R. C., 2004, AJ, 127, 1089
Lynch D.K., Russell R. W., Rudy R.J., Pearson R., Woodward C. E., 2007, IAU Circ. 8874
Marshall D. J., Robin A. C., Reyle C., Schultheis M., Picaud S., 2006, A&A, 453, 635
Munari, U.; Siviero A., Dallaporta S. et al., 2007a, CBET, 965.
Munari U., Orio M., Valentini M. et al., 2007b, CBET, 1010.
Munari U., Henden A., Banerjee D.P.K., Ashok N.M. , Righetti G.L., Dallaporta S. and Cetrulo G., 2014 (submitted)
Nakano S., Sakurai Y., Itagaki K., Koff R., 2007, IAU Circ. 8832
Naik S., Banerjee D. P. K., Ashok N. M., Das R. K., 2010, MNRAS, 404, 367
Naito H., Matsuda K., Yamaoka H., 2007, CBET 934
Poggiani R., 2008, NewA, 13, 557
Poggiani R., 2010, NewA, 15, 657
Raj A., Banerjee D. P. K., Ashok N. M., 2013, MNRAS, 433, 2657
Rudy R. J., Mazuk S., Puetter R. C., Hamann F., 2000, ApJ, 539, 166
Rudy R. J., Lynch D. K., Russell R. W., Woodward C. E., 2007, IAU Circ., 8884
Rudy R. J., Lynch D. K., Mazuk S., Venturini C. C., Puetter R. C., perry R. B., 2002, BAAS, 34, 1162
Shen K. J., Idan I., Bildsten L., 2009, ApJ, 705, 693
Shore S. N., 2013, A&A, 559, L7
Storey P. J., Hummer D. G., 1995, MNRAS, 292, 41
Strope R. J., Schaefer B. E., Henden A. A., 2010, AJ, 140, 34
Tanaka J., Nogami D., Fujii M, Ayani K., Kato T., Maehara H., Kiyota S., Nakajima K., 2011, PASJ, 63, 911.
Terzan A., 1970, POHP, 10, 167.
Wagner R. M., Depoy D. L., 1996, 467, 860
Walter F. M., Battisti A., Towers S. E., Bond H. E., Stringfellow G. S., 2012, PASP, 124, 1057.
Warner, B., 1995, *Cataclysmic Variable Stars. Cambridge Astrophysics Series*, Cambridge Univ. Press, Cambridge, New York, p. 260
Williams R.E., 1992, AJ, 104, 725
Williams R.E., 2012, AJ, 144, 98